

Infographic running myth: static stretching reduces injury risk in runners

James L N Alexander ¹, Christian J Barton,^{1,2} Richard W Willy³

It is a commonly held belief that static stretching plays an important role in improving running performance and decreasing injury risk.¹ As such static stretching, undertaken as part of a 'warm-up' prior to running, at the end of a run or as part of a strength training programme, is a common practice among runners of all levels. Static stretching involves

lengthening a muscle to the point at which a gentle tension is felt and remaining in this position, typically for a minimum of 30 seconds per stretch.²

Current research evidence definitively reports that this belief is, in fact, incorrect (figure 1).^{3,4}

Running places significantly high loads through the joints and soft tissues of the

body. As a result, runners are at a high risk of developing running-related overuse injuries such as patellofemoral joint pain, medial tibial stress syndrome, iliotibial band syndrome and Achilles tendinopathy.⁵ Overuse injuries most commonly occur when running loads (ie, frequency, intensity and/or duration) are progressed too rapidly and exceed the capacity of the tissue to adapt (ie, strengthen) to loading. This results in tissue overload and then injury. Other factors including muscle strength deficits, general health issues and injury history can also predispose a runner to injury.⁶

Research shows that stretching either in the short or long term has no impact on reducing the risk of overuse injuries experienced by endurance runners.^{3,4} Comparatively, an active warm-up, as described below, has been shown to improve running performance,⁷ although the benefit of an active warm-up for reducing injury risk in runners is still unclear.⁴

Static stretching is often recommended to enhance recovery and reduce the impact of delayed onset muscle soreness (DOMS). Endurance runners commonly experience DOMS after particularly fast or strenuous running (ie, running downhill). DOMS results in muscle pain, stiffness and fatigue, commonly in the thigh muscles, and can inhibit a runner's performance for several days.³ Unfortunately, stretching has not been shown to reduce the intensity or the duration of DOMS following exhaustive exercise.³

It is important to point out that regular static stretching after running will not have any detrimental effect on running performance.⁷

Static stretching will improve joint flexibility⁸ but these improvements have not been associated with benefits to recovery, performance or running economy in the long term.³ Runners should therefore consider other strategies that will assist them to prepare, perform and recover well from running, including the addition of a running specific strength-training programme into their training.⁹

A progressive warm-up prior to undertaking harder running efforts, training sessions or a race is recommended to optimise running performance.⁷ Performance gains are achieved via increased body



Figure 1 Infographic

temperature, metabolic changes, neural and psychological-related effects.⁷

As an alternative to passive stretching prior to running, a typical active warm-up for a long-distance runner could commence with 5–10 min of walking or easy jogging. If preparing for a fast running effort, training session or race, then incorporate 6–8 dynamic movement drills with particular focus on the lower limbs (ie, walking lunges, leg swings and A skips). Conclude the warm-up by completing three short efforts at goal running pace (ie, 3×100 m).⁷

Runners wishing to incorporate static stretching into their post running routine should do so, with the understanding that neither performance nor injury-prevention benefit will result.^{3 4}

¹Sports and Exercise Medicine Research Centre, La Trobe University, Melbourne, Victoria, Australia

²Department of Surgery, St Vincent's Hospital, University of Melbourne, Melbourne, Victoria, Australia

³Physical Therapy and Movement Science, University of Montana, Missoula, Montana, USA

Correspondence to James L N Alexander, La Trobe Sports and Exercise Research Centre, La Trobe University, Melbourne, VIC 3086, Australia; j.alexander@latrobe.edu.au

Twitter James L N Alexander @JamesA_15, Christian J Barton @DrChrisBarton and Richard W Willy @rwilly2003

Collaborators Katherine Parker.

Contributors JLNA and CJB: conceived the idea. JLNA: led development of the infographic and accompanying text. All authors contributed and approved the final infographic and accompanying text.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; internally peer reviewed.

© Author(s) (or their employer(s)) 2019. No commercial re-use. See rights and permissions. Published by BMJ.



To cite Alexander JLNA, Barton CJ, Willy RW. *Br J Sports Med* Epub ahead of print: [please include Day Month Year]. doi:10.1136/bjsports-2019-101169

Accepted 22 October 2019

Br J Sports Med 2019;0:1–2.

doi:10.1136/bjsports-2019-101169

ORCID iD

James L N Alexander <http://orcid.org/0000-0001-9474-6652>

REFERENCES

- 1 Saragiotto BT, Yamato TP, Lopes AD. What do recreational runners think about risk factors for running injuries? A descriptive study of their beliefs and opinions. *J Orthop Sports Phys Ther* 2014;44:733–8.
- 2 Woods K, Bishop P, Jones E. Warm-Up and stretching in the prevention of muscular injury. *Sports Med* 2007;37:1089–99.
- 3 Baxter C, Mc Naughton LR, Sparks A, et al. Impact of stretching on the performance and injury risk of long-distance runners. *Res Sports Med* 2017;25:78–90.
- 4 Yeung SS, Yeung EW, Gillespie LD, et al. Interventions for preventing lower limb soft-tissue running injuries. *Cochrane Database Syst Rev* 2011;56.
- 5 van Gent RN, Siem D, van Middelkoop M, et al. Incidence and determinants of lower extremity running injuries in long distance runners: a systematic review. *Br J Sports Med* 2007;41:469–80.
- 6 Saragiotto BT, Yamato TP, Hespanhol Junior LC, et al. What are the main risk factors for running-related injuries? *Sports Med* 2014;44:1153–63.
- 7 McGowan CJ, Pyne DB, Thompson KG, et al. Warm-Up strategies for sport and exercise: mechanisms and applications. *Sports Med* 2015;45:1523–46.
- 8 Bandy WD, Irion JM. The effect of time on static stretch on the flexibility of the hamstring muscles. *Phys Ther* 1994;74:845–50.
- 9 Blagrove RC, Howatson G, Hayes PR. Effects of strength training on the physiological determinants of middle- and long-distance running performance: a systematic review. *Sports Med* 2018;48:1117–49.